

HOMOLOGICAL DIMENSIONS OF THE AMALGAMATED DUPLICATION OF A RING ALONG A PURE IDEAL

MOHAMED CHHITI AND NAJIB MAHDOU

ABSTRACT. The aim of this paper is to study the classical global and weak dimensions of the amalgamated duplication of a ring R along a pure ideal I .

1. INTRODUCTION

Throughout this paper all rings are commutative with identity element and all modules are unitary.

Let R be a ring, and let M be an R -module. As usual we use $\text{pd}_R(M)$, $\text{id}_R(M)$ and $\text{fd}_R(M)$ to denote, respectively, the classical projective dimension, injective dimension and flat dimension of M . We use also $\text{gldim}(R)$ and $\text{wdim}(R)$ to denote, respectively, the classical global and weak dimension of R .

For a nonnegative integer n , an R -module M is n -presented if there is an exact sequence $F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_0 \rightarrow M \rightarrow 0$ in which each F_i is a finitely generated free R -module. In particular, “0-presented” means finitely generated and “1-presented” means finitely presented. Set $\lambda_R(M) = \{n/M \text{ is } n\text{-presented}\}$ except if M is not finitely generated. In this last case, we set $\lambda_R(M) = -1$. Note that $\lambda_R(M) \geq n$ is a way to express the fact that M is n -presented.

Given nonnegative integers n and d , a ring R is called an (n, d) -ring if every n -presented R -module has projective dimension $\leq d$, and R is called a weak (n, d) -ring if every n -presented cyclic R -module has projective dimension $\leq d$. For instance, the $(0, 1)$ -domains are the Dedekind domains, the $(1, 1)$ -domains are the Prüfer domains and the $(1, 0)$ -rings are the Von Neumann regular rings (see [1, 11, 12, 13, 14]). A commutative ring is called an n -Von Neumann regular ring if it is an $(n, 0)$ -ring. Thus, the 1-von Neumann regular rings are the von Neumann regular rings ([1, Theorem 1.3]).

The amalgamated duplication of a ring R along an ideal I is a ring that is defined as the following subring with unit element $(1, 1)$ of $R \times R$:

$$R \bowtie I = \{(r, r + i) / r \in R, i \in I\}$$

This construction has been studied, in the general case, and from the different point of view of pullbacks, by D’Anna and Fontana [6]. Also, in [5], they have considered the case of the amalgamated duplication of a ring, in not necessarily Noetherian setting, along a multiplicative canonical ideal in the sense of [10]. In [4], D’Anna has studied some properties of $R \bowtie I$, in order to construct reduced Gorenstein

2000 *Mathematics Subject Classification.* 13D05, 13D02.

Key words and phrases. Amalgamated duplication of a ring along an ideal, pure ideal, (n, d) -rings and weak (n, d) -rings.

rings associated to Cohen-Macaulay rings and has applied this construction to curve singularities. On the other hand, Maimani and Yassemi, in [16], have studied the diameter and girth of the zero-divisor of the ring $R \bowtie I$. Recently in [3], the authors study some homological properties of the rings $R \bowtie I$. Some references are [4, 5, 6, 16].

Let M be an R -module, the idealization $R \ltimes M$ (also called the trivial extension), introduced by Nagata in 1956 (cf [17]) is defined as the R -module $R \oplus M$ with multiplication defined by $(r, m)(s, n) = (rs, rn + sm)$ (see [7, 9, 11, 12]).

When $I^2 = 0$, the new construction $R \bowtie I$ coincides with the idealization $R \ltimes I$. One main difference of this construction, with respect to idealization is that the ring $R \bowtie I$ can be a reduced ring (and, in fact, it is always reduced if R is a domain). The first purpose of this paper is to study the classical global and weak dimension of the amalgamated duplication of a ring R along pure ideal R . Namely, we prove that if I is a pure ideal of R , then $\text{wdim}(R \bowtie I) = \text{wdim}(R)$. Also, we prove that if R is a coherent ring and I is a finitely generated pure ideal of R , then $R \bowtie I$ is an $(1, d)$ -ring provided the local ring R_M is an $(1, d)$ -ring for every maximal ideal M of R . Finally, we give several examples of rings which are not weak (n, d) -rings (and so not (n, d) -rings) for each positive integers n and d .

2. MAIN RESULTS

Let R be a commutative ring with identity element 1 and let I be an ideal of R . We define $R \bowtie I = \{(r, s)/r, s \in R, s - r \in I\}$. It is easy to check that $R \bowtie I$ is a subring with unit element $(1, 1)$, of $R \times R$ (with the usual componentwise operations) and that $R \bowtie I = \{(r, r + i)/r \in R, i \in I\}$.

It is easy to see that, if π_i ($i = 1, 2$) are the projections of $R \times R$ on R , then $\pi_i(R \bowtie I) = R$ and hence if $O_i = \ker(\pi_i \setminus R \bowtie I)$, then $R \bowtie I/O_i \cong R$. Moreover $O_1 = \{(0, i), i \in I\}$, $O_2 = \{(i, 0), i \in I\}$ and $O_1 \cap O_2 = (0)$.

Our first main result in this paper is given by the following Theorem:

Theorem 2.1. *Let R be a ring and I be a pure ideal of R . Then, $\text{wdim}(R \bowtie I) = \text{wdim}(R)$.*

To prove this Theorem we need some results.

Lemma 2.2. [4, Proposition 7] *Let R be a ring and let I be an ideal of R . Let P be a prime ideal of R and set:*

- $P_0 = \{(p, p + i)/p \in P, i \in I \cap P\}$,
- $P_1 = \{(p, p + i)/p \in P, i \in I\}$, and
- $P_2 = \{(p + i, p)/p \in P, i \in I\}$.

- (1) *If $I \subseteq P$, then $P_0 = P_1 = P_2$ and $(R \bowtie I)_{P_0} \cong R_P \bowtie I_P$.*
- (2) *If $I \not\subseteq P$, then $P_1 \neq P_2$, $P_1 \cap P_2 = P_0$ and $(R \bowtie I)_{P_1} \cong R_P \cong (R \bowtie I)_{P_2}$.*

Lemma 2.3. *Let I be a non-zero flat ideal of a ring R . For any R -module M we have:*

- (1) $fd_R(M) = fd_{R \bowtie I}(M \otimes_R (R \bowtie I))$.
- (2) $pd_R(M) = pd_{R \bowtie I}(M \otimes_R (R \bowtie I))$.

Proof. Note that the R -module $R \bowtie I$ is faithfully flat since I is flat. Firstly suppose that $fd_R(M) \leq n$ (resp., $pd_R(M) \leq n$) and pick an n -step flat (resp., projective) resolution of M over R as follows:

$$(*) \quad 0 \rightarrow F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_0 \rightarrow M \rightarrow 0.$$

Applying the functor $- \otimes_R R \bowtie I$ to $(*)$, we obtain the exact sequence of $(R \bowtie I)$ -modules:

$$0 \rightarrow F_n \otimes_R (R \bowtie I) \rightarrow F_{n-1} \otimes_R (R \bowtie I) \rightarrow \dots \rightarrow F_0 \otimes_R (R \bowtie I) \rightarrow M \otimes_R (R \bowtie I) \rightarrow 0$$

Thus $fd_{R \bowtie I}(M \otimes_R (R \bowtie I)) \leq n$ (resp., $pd_{R \bowtie I}(M \otimes_R (R \bowtie I)) \leq n$).

Conversely, suppose that $fd_{R \bowtie I}(M \otimes_R (R \bowtie I)) \leq n$ (resp., $pd_{R \bowtie I}(M \otimes_R (R \bowtie I)) \leq n$). Inspecting [2, page 118] and since $Tor_R^k(M, R \bowtie I) = 0$ for each $k \geq 1$, we conclude that for any R -module N and each $k \geq 1$ we have:

- (1): $Tor_R^k(M, N \otimes_R (R \bowtie I)) \cong Tor_{R \bowtie I}^k(M \otimes_R (R \bowtie I), N \otimes_R (R \bowtie I))$
- (2): $Ext_R^k(M, N \otimes_R (R \bowtie I)) \cong Ext_{R \bowtie I}^k(M \otimes_R (R \bowtie I), N \otimes_R (R \bowtie I))$

On the other hand $Tor_R^k(M, N)$ and $Ext_R^k(M, N)$ are direct summands of $Tor_R^k(M, N \otimes_R (R \bowtie I))$ and $Ext_R^k(M, N \otimes_R (R \bowtie I))$ respectively. Then, we conclude that $fd_R(M) \leq n$ (resp., $pd_R(M) \leq n$) and this finish the proof of this result. \square

One direct consequence of this Lemma is:

Corollary 2.4. *Let I be a non-zero flat ideal of a ring R . Then:*

- (1) $wdim(R) \leq wdim(R \bowtie I)$.
- (2) $gldim(R) \leq gldim(R \bowtie I)$.

Proof of Theorem 2.1. The inequality $wdim(R) \leq wdim(R \bowtie I)$ holds directly from Corollary 2.4 since I is pure and then flat. So, only the other inequality need a proof.

Using [7, Theorem 1.3.14] we have:

$$(\Upsilon) \quad wdim(R \bowtie I) = \sup\{wdim((R \bowtie I)_M) \mid M \text{ is a maximal ideal of } R \bowtie I\}.$$

Let M be an arbitrary maximal ideal of $R \bowtie I$ and set $m := M \cap R$. Then necessarily $M \in \{M_1, M_2\}$ where $M_1 = \{(r, r + i)/r \in m, i \in I\}$ and $M_2 = \{(r + i, r)/r \in m, i \in I\}$ (by [6, Theorem 3.5]). On the other hand, $I_m \in \{0, R_m\}$ since I is pure and m is maximal in R (by [7, Theorem 1.2.15]). Then, testing all cases of Lemma 2.3, we resume two cases;

- (1) $(R \bowtie I)_M \cong R_m$ if $I_m = 0$ or $I \not\subseteq m$.
- (2) $(R \bowtie I)_M \cong R_m \times R_m$ if $I_m = R_m$ or $I \subseteq m$.

Hence, we have $wdim((R \bowtie I)_M) = wdim(R_m) \leq wdim(R)$. So, the desired inequality follows from the equality (Υ) . \square

Corollary 2.5. *Let I be a finitely generated pure ideal of a ring R . Then R is a semihereditary ring if, and only if, $R \bowtie I$ is a semihereditary ring.*

Proof. Follows immediately from Theorem 2.1 and [3, Theorem 3.1]. \square

Recall that a ring R is called Gaussian if $c(fg) = c(f)c(g)$ for every polynomials $f, g \in R[X]$, where $c(f)$ is the content of f , that is, the ideal of R generated by the coefficients of f . See for instance [8].

Corollary 2.6. *Let R be a reduced ring and let I be a pure ideal of R . Then R is Gaussian if, and only if, $R \bowtie I$ is Gaussian.*

Proof. Follows immediately from Theorem 2.1 , [8, Theorem 2.2] and [6, Theorem 3.5(a)(vi)]. \square

By the fact that every ideal over a Von Neumann regular ring is pure, we conclude from Theorem 2.1 the following Corollary which have already proved in [3] with different methods.

Corollary 2.7. *Let R be a ring and let I be an ideal of R . If R is a Von Neumann regular ring, then so is $R \bowtie I$.*

If the ring R is Noetherian the global and weak dimensions coincide. Hence, Theorem 2.1 can be writing as follows:

Corollary 2.8. *If I is a pure ideal of a Noetherian ring R , then $\text{gdim}(R \bowtie I) = \text{gdim}(R)$.*

A simple example of Theorem 2.1 is given by introducing the notion of the trace of modules. Recall that if M is an R -module, the trace of M , $\text{Tr}(M)$, is the sum of all images of morphisms $M \rightarrow R_R$ (see [15]). Clearly $\text{Tr}(M)$ is an ideal of R .

Example 2.9. If M is a projective module over a ring R , then $\text{wdim}(R \bowtie \text{Tr}(M)) = \text{wdim}(R)$.

Proof. Clear since $\text{Tr}(M)$ is a pure ideal whenever M is projective (by [19, pp. 269-270]). \square

Now, we study the transfer of an $(1, d)$ -property.

Theorem 2.10. *Let R be a coherent ring such that for every maximal ideal m of R the local ring R_m is an $(1, d)$ -ring, and let I be a finitely generated pure ideal of R . Then $R \bowtie I$ is an $(1, d)$ -ring.*

Proof. Using [1, Theorem 3.2] and [3, Theorem 3.1], we have to prove that for any maximal ideal M of $R \bowtie I$, the ring $(R \bowtie I)_M$ is an $(1, d)$ -ring. So, let M be such ideal and set $m := M \cap R$. From the proof of Theorem 2.1, we have two possible cases:

- (1) $(R \bowtie I)_M \cong R_m$ if $I_m = 0$ or $I \not\subseteq m$.
- (2) $(R \bowtie I)_M \cong R_m \times R_m$ if $I_m = R_m$ or $I \subseteq m$.

So, by the hypothesis conditions, $(R \bowtie I)_M$ is an $(1, d)$ -ring since R_m is it, as desired. \square

By the fact that every ideal over a semisimple ring is pure we conclude from Theorem 2.10 the following Corollary.

Corollary 2.11. *Let R be a ring and let I be an ideal of R . If R is a semisimple ring, then so is $R \bowtie I$.*

Now, we give a wide class of rings which are not weak (n, d) -rings (and so not (n, d) -rings) for each positive integers n and d .

Theorem 2.12. *Let R be a ring and let I be a proper ideal of R which satisfies the following condition:*

- (1) R_m is a domain for every maximal ideal m of R .
- (2) I_m is a principal proper ideal of R_m for every maximal ideal m of R .

Then, $\text{wdim}(R \bowtie I) (= \text{gldim}(R \bowtie I)) = \infty$.

Proof. Let m be a maximal ideal of R such that $I \subseteq m \subsetneq R$. By Lemma 2.3, $R_m \bowtie I_m = (R \bowtie I)_M$ where $M = \{(p, p+i) | p \in m, i \in I\}$. From [3, Theorem 2.13] and by the hypothesis conditions, we have $\text{wdim}(R \bowtie I)_M = \text{wdim}(R_m \bowtie I_m) = \infty$. Then, the desired result follows from [7, Theorem 1.3.14]. \square

The following example shows that the condition " I_m is a principal proper ideal of R_m for every maximal ideal m in R " is necessary in Theorem 2.12.

Example 2.13. Let R be a Von Neumann regular ring and let I be a proper ideal of R . Then $\text{wdim}(R \bowtie I) = 0$ since $(R \bowtie I)$ is a Von Neumann regular ring, and I_m is not a proper ideal of R_m since R_m is a field.

REFERENCES

1. D. L. Costa; *Parameterising families of non-Noetherian rings*, Comm. Algebra, 22(1994), 3997-4011
2. H. Cartan and S. Eilenberg; *Homological Algebra*, Princeton Univ. Press. Princeton (1956).
3. M. Chhiti and N. Mahdou; *Some homological properties of an amalgamated duplication of a ring along an ideal*, Submitted for publication. Available from math.AC/0903.2240 V1 12 mar 2009
4. M. D'Anna; *A construction of Gorenstein rings*, J. Algebra **306** (2006), no. 2, 507-519.
5. M. D'Anna and M. Fontana; *The amalgamated duplication of a ring along a multiplicative-canonical ideal*, Ark. Mat. **45** (2007), no. 2, 241-252.
6. M. D'Anna and M. Fontana; *An amalgamated duplication of a ring along an ideal: the basic properties*. J. Algebra Appl. **6** (2007), no. 3, 443-459.
7. S. Glaz; *Commutative Coherent Rings*, Springer-Verlag, Lecture Notes in Mathematics, 1371 (1989).
8. S. Glaz; *The weak Dimensions of Gaussian rings*, Proc.Amer. Maths. Soc. 133 (2005), 2507-2513.
9. J. A. Huckaba; *Commutative Coherent Rings with Zero Divisors*. Marcel Dekker, New York Basel, (1988).
10. W. Heinzer, J. Huckaba and I. Papick; *m-canonical ideals in integral domains*, Comm.Algebra 26(1998), 3021-3043.
11. S. Kabbaj and N. Mahdou; *Trivial Extensions Defined by coherent-like condition*, Comm.Algebra 32 (10) (2004), 3937-3953
12. S. Kabbaj and N. Mahdou; *Trivial extensions of local rings and a conjecture of Costa*, Lecture Notes in Pure and Appl. Math., Vol.231, Marcel Dekker, New York, (2003), 301-312.
13. N. Mahdou; *On Costa's conjecture*, Comm. Algebra, 29 (7) (2001), 2775-2785.
14. N. Mahdou; *On 2-Von Neumann regular rings*, Comm. Algebra 33 (10) (2005), 3489-3496.
15. W.M. McGovern, G. Puninski, P. Rothmaler; *When every projective module is a direct sum of finitely generated modules*. J. Algebra 315 (2007). 454-481.
16. H. R. Maimani and S. Yassemi; *Zero-divisor graphs of amalgamated duplication of a ring along an ideal*. J. Pure Appl. Algebra **212** (1) (2008), 168-174.
17. M. Nagata; *Local Rings*. Interscience, New york, (1962).
18. G. Puninski and P. Rothmaler, *When every finitely generated flat module is projective*. J. Algebra 277 (2004), 542-558.
19. W. V. Vasconcelas; *Finiteness in projective ideals*. J. Algebra 25(1973). 269-278.

MOHAMED CHHITI, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY OF FEZ, BOX 2202, UNIVERSITY S.M. BEN ABDELLAH FEZ, MOROCCO., CHHITI.MED@HOTMAIL.COM

NAJIB MAHDOU, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY OF FEZ, BOX 2202, UNIVERSITY S.M. BEN ABDELLAH FEZ, MOROCCO., MAHDOU@HOTMAIL.COM